

Exercise 7: Strain

06.12.2024 - 09.12.2024

Question 1
 Consider the following displacement field,

$$\mathbf{u}(x, y, z) = k \begin{bmatrix} 2x + y^2 \\ x^2 - 3y^2 \\ 0 \end{bmatrix},$$

where k is a nonzero constant. Calculate the strain tensor ε !

Solution: The strain tensor ε was defined in the lecture by

$$\varepsilon = \frac{1}{2} \left(\vec{\nabla} \vec{u} + (\vec{\nabla} \vec{u})^T \right) \quad \text{or in index notation} \quad \varepsilon_{ij} = \frac{1}{2} \left(\frac{\partial u_i}{\partial r_j} + \frac{\partial u_j}{\partial r_i} \right)$$

With this formulas you are able to compute each component of the strain tensor


$$\begin{aligned} \varepsilon_{xx} &= \frac{1}{2} \left(\frac{\partial u_x}{\partial x} + \frac{\partial u_x}{\partial x} \right) = \frac{\partial(2x + y^2)}{\partial x} = 2k \\ \varepsilon_{xy} &= \frac{1}{2} \left(\frac{\partial u_x}{\partial y} + \frac{\partial u_y}{\partial x} \right) = \frac{1}{2} \left(\frac{\partial(2x + y^2)}{\partial y} + \frac{\partial(x^2 - 3y^2)}{\partial x} \right) = \frac{1}{2} (2ky + 2kx) = k(x + y) \\ \varepsilon_{yy} &= \frac{1}{2} \left(\frac{\partial u_y}{\partial y} + \frac{\partial u_y}{\partial y} \right) = \frac{\partial(x^2 - 3y^2)}{\partial y} = -6ky \\ &\dots \end{aligned}$$

By using the symmetry of ε you only have to compute 6 entries and should find the following result

$$\varepsilon = k \begin{bmatrix} 2 & x + y & 0 \\ x + y & -6y & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

Note that all components of ε involving the z -direction are zero. This situation is called *plane strain*.

Question 2
 A solid bar with dimensions $l \times w \times w$ (see below) is stretched along its length to a final length al . The volume of the bar does not change during deformation. Calculate the displacement vector \mathbf{u} and the strain tensor ε !



Solution: After stretching, the bar has a new width \hat{w} . However, the volume is conserved, therefore

$$\hat{w}^2 l a = w^2 l \implies \hat{w} = \frac{w}{\sqrt{a}}.$$

We assume the bar deforms homogeneously. The displacement along the x -direction increases linearly from zero at $x = 0$ to $(a - 1)l$ at $x = l$. Similarly, the displacement in y -direction increases linearly from zero at $y = 0$ to some maximum value at $y = w/2$. The displacement in z -direction increases linearly from zero at $z = 0$ to some maximum value at $z = w/2$. The bar retains its square cross-section, i.e. the y -displacement does not depend on x and z , and the z -displacement does not depend on y and x . Therefore, the displacement vector can be written as

$$\mathbf{u} = \begin{bmatrix} (a - 1)x \\ Ay + B \\ Cz + D \end{bmatrix},$$

where A, B, C, D are constants that need to be determined by consideration of the boundary conditions. Since the y - and z - components are zero at $y = 0$ and $z = 0$, respectively, we see that $B = D = 0$. The bar retains its square cross-section, i.e. the y -displacement at $y = w/2$ must be equal to the z -displacement at $z = w/2$. Therefore $A = C$. Recall that the new width after deformation is $\hat{w} = w/\sqrt{a}$. For this reason, the y -displacement at $y = w/2$ must be equal to $(\hat{w} - w)/2 = (1/\sqrt{a} - 1)w/2$. It must also be equal to $Aw/2$. Therefore $A = (1/\sqrt{a} - 1)$. In conclusion, the displacement vector is

$$\mathbf{u} = \begin{bmatrix} (a - 1)x \\ \left(\frac{1}{\sqrt{a}} - 1\right)y \\ \left(\frac{1}{\sqrt{a}} - 1\right)z \end{bmatrix}.$$

Through differentiation, we obtain the strain tensor

$$\varepsilon = \begin{bmatrix} a - 1 & 0 & 0 \\ 0 & \frac{1}{\sqrt{a}} - 1 & 0 \\ 0 & 0 & \frac{1}{\sqrt{a}} - 1 \end{bmatrix}.$$

Question 3
 (Saint-Venant's compatibility conditions)

The strain tensor ε has six distinct components. However, these six components are computed from only three components of the displacement vector \mathbf{u} . Thus, if we want to solve for the components of \mathbf{u} given the component of ε , we have six equations for three unknowns. For this system of equations to have a solution, some of the strain components must be related. Show that they are by considering their second derivatives! For example, differentiate ε_{xx} twice with respect to y , ε_{yy} twice with respect to x and ε_{xy} with respect to x and y , and compare!

Solution: Recall the definition of the aforementioned strain components,

$$\begin{aligned} \varepsilon_{xx} &= \frac{\partial u_x}{\partial x}, \\ \varepsilon_{yy} &= \frac{\partial u_y}{\partial y}, \\ \varepsilon_{xy} &= \frac{1}{2} \left(\frac{\partial u_x}{\partial y} + \frac{\partial u_y}{\partial x} \right). \end{aligned}$$

Thus,

$$\frac{\partial^2 \varepsilon_{xx}}{\partial y^2} = \frac{\partial^3 u_x}{\partial x \partial y^2},$$

$$\frac{\partial^2 \varepsilon_{yy}}{\partial x^2} = \frac{\partial^3 u_y}{\partial y \partial x^2},$$

and

$$\frac{\partial^2 \varepsilon_{xy}}{\partial x \partial y} = \frac{1}{2} \frac{\partial^2}{\partial x \partial y} \left(\frac{\partial u_x}{\partial y} + \frac{\partial u_y}{\partial x} \right),$$

$$= \frac{1}{2} \left(\frac{\partial^3 u_x}{\partial y^2 \partial x} + \frac{\partial^3 u_y}{\partial x^2 \partial y} \right).$$

The order of differentiation in the last equation is immaterial, hence we see that

$$\frac{\partial^2 \varepsilon_{xx}}{\partial y^2} + \frac{\partial^2 \varepsilon_{yy}}{\partial x^2} = 2 \frac{\partial^2 \varepsilon_{xy}}{\partial x \partial y}.$$

Two more equations of this form can be obtained by repeating this procedure for ε_{yy} , ε_{zz} , and ε_{yz} , and for ε_{xx} , ε_{zz} , and ε_{xz} . This is tantamount to cyclic permutations of the indices, $x \rightarrow y$, $y \rightarrow z$, and $z \rightarrow x$.

Three more equations can be obtained by considering mixed derivatives of type

$$\frac{\partial^2 \varepsilon_{xx}}{\partial y \partial z} = \frac{\partial^3 u_x}{\partial x \partial y \partial z}.$$

Again, order of differentiation is immaterial, hence

$$\frac{\partial^3 u_x}{\partial x \partial y \partial z} = \frac{\partial^2}{\partial x \partial z} \left(\frac{\partial u_x}{\partial y} \right)$$

$$= \frac{\partial^2}{\partial x \partial z} \left(2\varepsilon_{xy} - \frac{\partial u_y}{\partial x} \right)$$

$$= \frac{\partial}{\partial x} \left(2 \frac{\partial \varepsilon_{xy}}{\partial z} - \frac{\partial^2 u_y}{\partial x \partial z} \right).$$

Similarly,

$$\frac{\partial^3 u_x}{\partial x \partial y \partial z} = \frac{\partial^2}{\partial x \partial y} \left(\frac{\partial u_x}{\partial z} \right)$$

$$= \frac{\partial^2}{\partial x \partial y} \left(2\varepsilon_{xz} - \frac{\partial u_z}{\partial x} \right)$$

$$= \frac{\partial}{\partial x} \left(2 \frac{\partial \varepsilon_{xz}}{\partial y} - \frac{\partial^2 u_z}{\partial x \partial y} \right).$$

Combing gives

$$2 \frac{\partial^3 u_x}{\partial x \partial y \partial z} = \frac{\partial}{\partial x} \left(2 \frac{\partial \varepsilon_{xy}}{\partial z} - \frac{\partial^2 u_y}{\partial x \partial z} \right) + \frac{\partial}{\partial x} \left(2 \frac{\partial \varepsilon_{xz}}{\partial y} - \frac{\partial^2 u_z}{\partial x \partial y} \right)$$

$$= \frac{\partial}{\partial x} \left(2 \frac{\partial \varepsilon_{xy}}{\partial z} + 2 \frac{\partial \varepsilon_{xz}}{\partial y} - \frac{\partial u_y}{\partial z} - \frac{\partial u_z}{\partial y} \right)$$

$$= 2 \frac{\partial}{\partial x} \left(\frac{\partial \varepsilon_{xy}}{\partial z} + \frac{\partial \varepsilon_{xz}}{\partial y} - \frac{\partial \varepsilon_{yz}}{\partial x} \right).$$

Finally,

$$\frac{\partial^2 \varepsilon_{xx}}{\partial y \partial z} = \frac{\partial}{\partial x} \left(\frac{\partial \varepsilon_{xy}}{\partial z} + \frac{\partial \varepsilon_{xz}}{\partial y} - \frac{\partial \varepsilon_{yz}}{\partial x} \right)$$

The other two equations of this type can be obtained by cyclic permutation of the indices, $x \rightarrow y, y \rightarrow z$, and $z \rightarrow x$. The six compatibility conditions are therefore

$$2 \frac{\partial^2 \varepsilon_{xy}}{\partial x \partial y} = \frac{\partial^2 \varepsilon_{xx}}{\partial y^2} + \frac{\partial^2 \varepsilon_{yy}}{\partial x^2} \quad (\text{a}),$$

$$2 \frac{\partial^2 \varepsilon_{yz}}{\partial y \partial z} = \frac{\partial^2 \varepsilon_{yy}}{\partial z^2} + \frac{\partial^2 \varepsilon_{zz}}{\partial y^2} \quad (\text{b}),$$

$$2 \frac{\partial^2 \varepsilon_{zx}}{\partial z \partial x} = \frac{\partial^2 \varepsilon_{zz}}{\partial x^2} + \frac{\partial^2 \varepsilon_{xx}}{\partial z^2} \quad (\text{c}),$$

$$\frac{\partial^2 \varepsilon_{xx}}{\partial y \partial z} = \frac{\partial}{\partial x} \left(\frac{\partial \varepsilon_{xy}}{\partial z} + \frac{\partial \varepsilon_{xz}}{\partial y} - \frac{\partial \varepsilon_{yz}}{\partial x} \right) \quad (\text{d}),$$

$$\frac{\partial^2 \varepsilon_{yy}}{\partial z \partial x} = \frac{\partial}{\partial y} \left(\frac{\partial \varepsilon_{yz}}{\partial x} + \frac{\partial \varepsilon_{yx}}{\partial z} - \frac{\partial \varepsilon_{zx}}{\partial y} \right) \quad (\text{e}),$$

$$\frac{\partial^2 \varepsilon_{zz}}{\partial x \partial y} = \frac{\partial}{\partial z} \left(\frac{\partial \varepsilon_{zx}}{\partial y} + \frac{\partial \varepsilon_{zy}}{\partial x} - \frac{\partial \varepsilon_{xy}}{\partial z} \right) \quad (\text{f}).$$

Question 4

In the first question, you computed the strain tensor ε for displacement field

$$\mathbf{u}(x, y, z) = k \begin{bmatrix} 2x + y^2 \\ x^2 - 3y^2 \\ 0 \end{bmatrix}.$$

Now show that ε fulfills the compatibility conditions!

Solution: Recall that

$$\varepsilon = kH \begin{bmatrix} 2 & x + y & 0 \\ x + y & -6y & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

Note that all components of ε involving the z -direction are zero (*plane strain*). All derivatives with respect to z are zero. Moreover, ε is linear in x and y . Therefore, all terms of the type $\partial^2(\dots)/\partial x^2$ and $\partial^2(\dots)/\partial y^2$ are zero. The only non zero term that is left over is $\frac{\partial^2 \varepsilon_{xy}}{\partial x \partial y}$. We can inspect this term in the first condition (a)

$$\begin{aligned} 2 \frac{\partial^2 \varepsilon_{xy}}{\partial x \partial y} &= 0 + 0 \\ \Leftrightarrow 2 \frac{\partial}{\partial y} \left(\frac{\partial \varepsilon_{xy}}{\partial x} \right) &= 0 \\ \Leftrightarrow 2 \frac{\partial}{\partial y} (1) &= 0 \\ \Leftrightarrow 0 &= 0 \quad \text{OK!} \end{aligned}$$

so the derivative $\frac{\partial^2 \varepsilon_{xy}}{\partial x \partial y} = 0$. Thus, we can immediately see that conditions (b)–(f) are fulfilled. Therefore the strains are compatible.